

LMDZ Single Column Model

- + what is it ?
- + why is it interesting ?
- + List of 1D cases
- + how to install and run it ?

M-P Lefebvre and LMDZ team


Use of 1D cases



Test case, field campaign experiment




Observation




Explicit simulations, Grid cell, 20-100 m

Evaluation



Climate model, parameterizations, « single-column » mode


« Large scale » conditions imposed



Courtesy F.Hourdin


3D is a collection of many “single column models”, covering earth and interaction with each other through a set of rules known as “large scale dynamics”.

In a 1D model, there is no dynamics. We use observations or model output or idealized forcing to impose forcing at the boundary of the column.



LMDZ model in 1D mode

- We impose large scale conditions.
- Duration of the case varies from few hours to few months
- We study parameterizations in a given environment.




Why use SCMs ?

- + **simplicity**: technical and understanding, usable on any laptop
- + it's a useful tool for **parameterization development**: shallow convection, deep convection, transition from stratocumulus to cumulus, stable boundary layer, radiation...
- + we can **compare results to observations or to explicit simulations** (CRM, LES)
- + then we go back to GCM and test new parameterizations ...
- + hierarchy of models: SCM, CRM, AGCM, GCM ...

shallow convection

- Arm_cu (diurnal cycle of shallow cumulus over land)
- Rico (shallow precipitating cumulus over sea)
- Ayotte (convective boundary layer)




Arm_cu case - Cloud fraction

Stratocumulus and transition to cumulus


- Sandu (transition case with 3 options according to variation of SST)
- Fire (diurnal cycle of stratocumulus)

Fire case:
Cloud fraction (%)



Top: standard version
Bad representation because not
Enough entrainment at the cloud top

Bottom: version developed by A.Jam






Deep convection:

Over ocean:


- Toga
- case_e (part of Toga)
- TWPICE

TWPICE Case





LES




LMDZ_AR4_L39

AMMA case:
Thetal tendencies due to all schemes (K/j)

Deep convection:

Over land:


- Hapex
- AMMA
- Idealized case:
eq_rad_conv (RCE)



LMDZ_NP_L70

How to proceed ?


- + install LMDZ 3D with **install.sh**
- + install LMDZ 1D with **instal1d.sh**.
 - * creates 1D directory
 - * compiles
 - * runs 6 cases (arm_cu, rico, sandufast, fire, twpice, amma) with 2 different physics and show cloud cover + precipitation plot



What can you do in each case directory ?

- + **compile & run** with run.sh: choose case, physics and level number
- + Look at profils and forcings (ascii files or netcdf files)
- + Modify *def files
- + Read readme file
- + **get results** in netcdf files (hist*.nc)
- + **compare** to LES results in resul_LES

Cases to run with different physical packages



How to run a case or compile after modifications : With run.sh

Which case(s) ?

```
listecas="dice ihop arm_cu rico sandufast sanduref sanduslow fire  
toga ayotte twpice case_e amma" # testes
```

```
listecas="arm_cu rico sandufast fire twpice amma"
```

```
listecas="amma"
```

Which physics ?

```
listedef="SP NPv3.1 NPv3.2 NPv4.12 NPv5.00 NPv5.10"
```

```
listedef="NPv5.00 "
```

```
listedef="NPv3.2 NPv5.00 "
```

Number of levels ?

```
case $DEF in
```

```
SP|NPV3.1|NPv3.2) L=39 ;;
```

```
NPv4.12) L=59 ;;
```

```
*) L=79
```

```
esac
```

Where are the results ?

In LMDZtesting/1D/RESU/NPv3.2L39/amma

lrwxrwxrwx 1 ...	36	3 déc.	11:38	amma.nc	← forcings
-rw-r--r-- 1 ...	285452	3 déc.	11:38	histhf.nc	← Result files
-rw-r--r-- 1 ...	279088	3 déc.	11:38	hourly.nc	→
-rw-r--r-- 1 ...	15292	3 déc.	11:38	limit.nc	
-rw-r--r-- 1 ...	652	3 déc.	11:38	lmdz1d.def	
-rw-r--r-- 1 ...	4247	3 déc.	11:38	config.def	* def files
-rw-r--r-- 1 ...	85	3 déc.	11:38	gcm.def	
-rw-r----- 1 ...	3765	3 déc.	11:38	physiq.def	
-rwxr-xr-- 1 ...	692	3 déc.	11:38	run.def	
-rw-r--r-- 1 ...	42	3 déc.	11:38	traceur.def	
-rw-r--r-- 1 ...	7728	3 déc.	11:38	paramLMDZ_phy.nc	
-rw-r--r-- 1 ...	16532	3 déc.	11:38	startphy.nc	

CAUTION !

Don't modify *def files in ~LMDZtesting/1D/RESU directory !!

Here you have only copies of the files.



The « original » files are either under ~LMDZtesting/CAS or ~LMDZtesting/INPUT

Eurocs Cumulus

Rico



LES

Z (m)






NPv3

Z (m)



SP

Z (m)





In LMD, we used these cases to develop New Physics version.

For 2 cases, Arm_cu and Rico, we compare results of « standard physics » (CMIP3), « new physics » (CMIP5) and LES model.

Shade= cloud cover
Contour= specific humidity (g/kg)

Rico case :

Sensitivity Tests to vertical
discretization and time step



Recent improvements:

+ 1D with tracers

+ 1D coupled with soil model Orchidee:



DICE case : characterize boundary layer
In the site of SGP during 3 days/night



GABLS4 case : interaction of a very
stable boundary layer with a snow surface